

COSII Marker C2_At3g52220, 10.00cM

Christopher Martin and Douglas Maxwell
University of Wisconsin-Madison: June 19, 2006

Fig.1: Map of the top of Chr. 11. (Modified from Solanaceae Genomics Network, 2006).




Table 1: Primers from C2_At3g52220 on Chr. 11

Primer Name	Sequence (5'-3')
DM11 - F2	TGCTCGGGTGGATGGTCTTGG
DM11 - R2	TGATGGTGAACCTGGTTCTTCCC




Fig. 2: Agarose gel of the PCR reactions with the DM11F2/DM11R2 primers from Chr. 11. lane 1, Heinz 1706; lane 2, Gh13; lane 3, Gc9; lane 4, LA1777; lane 5, LA2779; lane 6, water; lane 7, 100-bp ladder. Arrow marks the 700-bp fragment.

Background: The purpose of this project was to locate molecular markers for disease resistance in tomato. To accomplish this goal, primers were obtained from the Solanaceae Genomics Network (SGN) website (Solanaceae Genomics Network, 2006), and used with five different tomato lines.

We used the tomato breeding lines, Gh13 and Gc9 which are resistant to the bipartite begomoviruses in Guatemala (Mejía *et al.*, 2004; Nakhla *et al.*, 2004). Gh13 is the F7 generation and is a homogeneous breeding line with resistance derived from *L. hirsutum*. Gc9 is at least an F8 breeding line with resistance genes introgressed from *L. chilense* by J. W. Scott (Scott *et al.*, 1995). LA1777 is the *L. hirsutum* parent, and is thought to be the source of the introgression in Gh13. LA2779 is the *L. chilense* parent and is thought to be the source of the introgression in Gc9 (Maxwell, D., pers. com.)

As a control, we used the breeding line Heinz 1706. Heinz 1706 is the tomato cultivar being sequenced in an international sequencing project (Budiman *et al.*, 2000; Ozminowski, 2004), and is susceptible to geminiviruses (Hapidat, M., pers. com.). The susceptibility of Heinz 1706 to geminiviruses was confirmed through testing with *Tomato Yellow Leaf Curl Virus*, which is a begomovirus (Maxwell, D., pers. com.).

The begomovirus resistant lines, Gh13 and Gc9 were supplied by Dr. L. Mejía, Universidad de San Carlos, Guatemala City. The susceptible line, Heinz 1706, was supplied by Dr. R. Ozminowski, Heinz Seed Co., Stockton, CA.

Polymerase Chain Reaction (PCR): PCR fragments from each set of primers, for each of the five genotypes, were obtained using methods developed in the Maxwell lab (Czosnek *et al.*, 2004). PCR parameters were for 50- μ l reactions containing: 5- μ l 2.5mM deoxynucleotide triphosphates (dNTPs), 5- μ l 10X buffer, 5- μ l 25 mM MgCl₂, 0.2- μ l *Taq* DNA polymerase, 5- μ l each forward and reverse sense primer at 10 μ M, 5-7 μ l of DNA extract, and H₂O. Some PCR reactions were run with 25- μ l reactions. When this was the case, the concentrations of all chemicals were exactly half of what appeared in the 50- μ l reactions. PCR cycle parameters for fragment amplification were as follows: denaturation at 94°C for 3 min, then 35

cycles at 94°C for 30 sec each, annealing at 53°C for 1 min, and extension at 72°C for 1 min. These cycles were followed by a reaction at 72°C for 10 min, and then the reaction was held at 4°C. PCR reactions were performed in the MJ DNA Engine PT200 Thermocycler™ (MJ Research Inc., Waltham, MA).

The PCR-amplified DNA was run on an electrophoresis gel of 1.5% Seakem LE™ agarose (BioWhittaker Molecular Applications Rockland, ME) in 0.5X TBE buffer, stained with ethidium bromide, and visualized with a Kodak Gel Logic 200 Imaging System.

DM11F2-R2 Results: The DM11F2/DM11R2 primer pair was chosen from the list of COSII primers on the SGN website (Solanaceae Genomics Network, 2006). The primer combination produced three distinct patterns. Heinz 1706, Gh13, and Gc9 all produced four bands. Strong bands were at 600bp, 500bp and 350bp. A weak band appeared at 200bp. LA1777 and LA2779 each had unique banding patterns. LA1777 produced three bands with one at 600, 450, and 400bp respectively. LA2779 also produced three bands with one at 600, 550, and 400bp respectively. Because these primers produced multiple bands, the PCR fragments were not sequenced.

Thus, the breeding lines had significantly different banding patterns from the species. This indicates that none of the breeding lines have an introgression in this region. These primers may be useful in tracking introgressions from LA2779 or LA1777 in this region. The different banding patterns would allow for a quick and easy analysis of the introgression through PCR.

Table 2: Results of the PCR with the DM11F2/DM11R2 primer pair.

Plant Line	Result
Heinz, Gh13, and Gc9	Four bands (strong bands at 600,500,350bp and a weak band at 200bp)
LA1777	Three strong bands (600, 450, and 400bp)
LA2779	Three strong bands (600bp, 550bp, and 400bp)

References

2006. "COS II Markers." Solanaceae Genomics Network. Cornell University. <http://www.sgn.cornell.edu/markers/cosii_markers.pl> (July 19, 2006).
- Budiman, MA., Mao, L., Wood, TC., and Wing, RA. 2000. A deep-coverage tomato BAC library and prospects toward development of an STC framework for genome sequencing. *Genome Res.* 10:129-136.
- Czosnek, H., Vidavski, F., Mejía, L., Lapidot, M., Maxwell, D., and Havey, M. 2004. "Molecular Marker-Assisted Breeding for Resistance to Whitefly-Transmitted Geminiviruses Infecting Tomato in Guatemala." *Achievements for CDR Grant 2004*. <<http://www.plantpath.wisc.edu/InVirLab/CDR-Grant04.htm>> (October 5, 2004).
- Mejía, L., Teni, R.E., Vidavski, F., Czosnek, H., Lapidot, M., Nakhla, M.K., and Maxwell D.P. 2004. Evaluation of tomato germplasm and selection of breeding lines for resistance to begomoviruses in Guatemala. *Acta Hort.* (in press).
- Nakhla, M., Sorenson, A., Mejía, L., Ramírez, P., Karkashian, J.P., and Maxwell, D., "Molecular Characterization of Tomato-Infecting Begomoviruses in Central America and Development of DNA-Based Detection Methods." *International Plant Virology Laboratory*. <<http://www.plantpath.wisc.edu/invirlab/docs/Beg-CA-Final.htm>> (October 5, 2004).
- Omnikowski, R. 2004. Pedigree of variety Heinz 1706. Report of the Tomato Genetics Cooperative 54: 27.
- Scott, J.W., Stevens, M.R., Barten, J.H.M., Thome, C.R., Polston, J.E., Schuster, D.J. and Serra, C.A. 1995. Introgression of resistance to whitefly-transmitted geminiviruses from *Lycopersicon chilense* to tomato. *Taxonomy, Biology, Damage Control and Management*, Ed. by D. Gerling and R.T. Mayer, Intercept Ltd., Andover, UK. p. 357-367.